


Shelter Placement for Patients with COVID-related Risk



^{*} Keep in mind that an individual may have recently tested positive at another hospital. If available, consider checking CRISP, or ask about the first positive test in the last 90 days. https://www.cdc.gov/coronavirus/2019-ncov/hcp/duration-isolation.html

https://www.cdc.gov/coronavirus/2019-ncov/more/scientific-brief-options-to-reduce-quarantine.html

<u>Lord Baltimore TRI Center</u>: https://coronavirus.baltimorecity.gov/isolation-assistance-tri-center-lord-baltimore-hotel-0

<u>Baltimore Convention Center Field Hospital</u>: https://www.umms.org/coronavirus/fighting-covid/expanded-clinical-locations/baltimore-convention-center/healthcare-providers
Mayor's Office of Homeless Services: https://homeless.baltimorecity.gov/get-help-now

^{**}COVID-like illness is defined as 1) One or more of the following: cough, shortness of breath, or new loss of taste or smell, OR 2) Two or more of the following: fever, chills, muscle aches, headache, sore throat, nausea or vomiting, diarrhea, fatigue, congestion, or runny nose.

^{***} CDC provides options to shorten quarantine. Baltimore City Health Department contact tracers use 10 days from last exposure if continuously asymptomatic. Lord Baltimore TRI Center provides additional testing prior to ending the 10-day quarantine in order to further reduce the chance of transmission upon return to a congregate setting such as a homeless shelter.